


Hydrocarbons-2

Lec. Haider Abdulkareem AlMashhadani


Lecture 11

2.2. Unsaturated H.C

- **Alkanes** are called saturated hydrocarbons, because they contain the maximum number of hydrogen atoms per carbon. In contrast, **alkenes and alkynes** are called unsaturated hydrocarbons.
- **Unsaturated hydrocarbons** are compounds that contain fewer than the maximum number of hydrogen atoms per carbon (or contain double or triple bond).
- **Alkenes and alkynes** are two families of organic molecules that contain multiple bonds.
- **Alkenes** are compounds that contain a carbon–carbon double bond.
- The general molecular formula of an alkene is C_nH_{2n} , so an alkene has **two fewer** hydrogens than an alkane.



- ❑ **Alkynes** are compounds that contain a carbon–carbon triple bond.
- ❑ The general molecular formula for an alkyne is C_nH_{2n-2} , so an alkyne has **four fewer** hydrogens than an acyclic alkane.




- ❑ Because alkenes and alkynes are composed of nonpolar carbon–carbon and carbon–hydrogen bonds, their physical properties are similar to other hydrocarbons. Like alkanes.

Alkenes and alkynes have low melting points and boiling points and are insoluble in water.


3. Isomerism and Stereoisomers

- **Isomers** are two different compounds with the same molecular formula.




butane


=



4 C's in a row




straight-chain alkane




isobutane

=





3 C's with a one-carbon branch





branched-chain alkane

- ❑ They are two major classes of isomers:
 - ✓ **Constitutional isomers:** differ in the way the atoms are connected to each other.
 - ✓ **Stereoisomer:** are isomers that differ only in the 3D arrangement of atoms.
- ❑ Constitutional isomers like butane and isobutane belong to the same family of compounds: they are both alkanes.
- ❑ Ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) and dimethyl ether (CH_3OCH_3) are constitutional isomers with different functional groups: $\text{CH}_3\text{CH}_2\text{OH}$ is an alcohol and CH_3OCH_3 is an ether.



ethanol




dimethyl ether

□ Stereoisomers:

- 2-Butene illustrates another important aspect about alkenes. There is restricted rotation around the carbon atoms of a double bond. As a result, the groups on one side of the double bond cannot rotate to the other side.
- With 2-butene, there are two ways to arrange the atoms on the double bond. The two CH₃ groups can be on the same side of the double bond or they can be on opposite sides of the double bond. These molecules are different compounds with the same molecular formula; that is, they are isomers.


- When the two CH₃ groups are on the same side of the double bond, the compound is called the *cis isomer*.
- When the two CH₃ groups are on opposite sides of the double bond, the compound is called the *trans isomer*.

General structure




2-butene

Two possible arrangements



two CH₃ groups
on the same side

cis isomer



two CH₃ groups
on opposite sides

trans isomer

SAMPLE PROBLEM 11.3


Draw *cis*- and *trans*-3-hexene.

Analysis

First, use the parent name to draw the carbon skeleton, and place the double bond at the correct carbon; 3-hexene indicates a 6 C chain with the double bond beginning at C3. Then use the definitions of *cis* and *trans* to draw the isomers.


Solution

Each C of the double bond is bonded to a CH_3CH_2 group and a hydrogen. A *cis* isomer has the CH_3CH_2 groups bonded to the same side of the double bond. A *trans* isomer has the two CH_3CH_2 groups bonded to the opposite sides of the double bond.



PROBLEM

Label each pair of alkenes as constitutional isomers or stereoisomers.




4. Reactions of Alkenes

- Most families of organic compounds undergo a characteristic type of reaction. Alkenes undergo addition reactions. In an **addition reaction**, new groups X and Y are added to a starting material. One bond of the double bond is broken and two new single bonds are formed.

➤ Addition reaction is a reaction in which elements are added to a compound.


- **Why does addition occur?**
- A double bond is composed of one strong bond and one weak bond. In an addition reaction, the weak bond is broken and two new strong single bonds are formed.

Addition



4.1. Addition of Hydrogen—Hydrogenation


- Hydrogenation is the addition of hydrogen (H_2) to an alkene. Two bonds are broken: one bond of the carbon–carbon double bond and the H-H bond—and two new C-H bonds are formed.



4.2. Addition of Water—Hydration


- Hydration is the addition of water to an alkene. Two bonds are broken—one bond of the carbon–carbon double bond and the H–OH bond—and new C–H and C–OH bonds are formed.

■



- **Hydration** occurs only if a **strong acid** such as **H₂SO₄** is added to the reaction mixture. The product of hydration is an *alcohol*.

- There is one important difference in this addition reaction compared to the addition of H_2 . In this case, addition puts different groups-H and OH-on the two carbons of the double bond. As a result, H_2O can add to the double bond to give two constitutional isomers when an unsymmetrical alkene is used as starting material.




only product

- This is specific example of a general trend called **Markovnikov's rule**

➤ In the addition of H_2O to an unsymmetrical alkene, the H atom bonds to the less substituted carbon atom—that is, the carbon that has more H's to begin with.


5. Aromatic Compounds

- **Aromatic compounds** represent another example of unsaturated hydrocarbons. Aromatic compounds were originally named because many simple compounds in this family have characteristic odors. Today, the word **aromatic refers to compounds that contain a benzene ring, or rings that react in a similar fashion to benzene.**
- **Benzene**, the simplest and most widely known aromatic compound, contains a six-membered ring and three double bonds. Since each carbon of the ring is also bonded to a hydrogen atom, the molecular formula for benzene is C_6H_6 .




benzene
 C_6H_6

=



=




planar molecule


HEALTH NOTE



Tamoxifen, a potent anticancer drug sold under the trade name of Novaldex, contains three benzene rings.



The pain reliever acetaminophen (trade name Tylenol) contains a para-disubstituted benzene ring.



acetaminophen
(Trade name: Tylenol)

6.1. Saturated and Unsaturated Fatty Acids

(FOCUS ON HEALTH & MEDICINE)

- Naturally occurring animal fats and vegetable oils are formed from fatty acids.
- **Fatty acids** are carboxylic acids (RCOOH) with long carbon chains of 12–20 carbon atoms. Because a fatty acid has many nonpolar C-C and C-H bonds and few polar bonds, fatty acids are **insoluble in water**. There are two types of fatty acids.

- **Saturated fatty** acids have no double bonds in their long hydrocarbon chains.
- **Unsaturated fatty** acids have one or more double bonds in their long hydrocarbon chains.

Generally, double bonds in naturally occurring fatty acids are **cis**.


As the number of double bonds in the fatty acid *increases*, the melting point *decreases*.

- **Fats** are solids at room temperature. Fats are generally formed from fatty acids having few double bonds. (from animal sources)
- **Oils** are liquids at room temperature. Oils are generally formed from fatty acids having a larger number of double bonds. (from vegetable sources)


6.2. Sunscreens

(FOCUS ON HEALTH & MEDICINE)

- All commercially available sunscreens contain a benzene ring. A sunscreen absorbs ultraviolet radiation and thus shields the skin for a time from its harmful effects. Two sunscreens that have been used for this purpose are **p-aminobenzoic acid (PABA)** and **Padimate O**.



p-aminobenzoic acid
(PABA)



Padimate O




Commercial sunscreens are given an **SPF** rating (sun protection factor), according to the amount of sunscreen present. The higher the number, the greater the protection.


6.3. Phenols as Antioxidants

(FOCUS ON HEALTH & MEDICINE)

- A wide variety of **phenols**, compounds that contain a **hydroxyl group** bonded to a **benzene ring**, occur in nature. **Vanillin** from the vanilla bean is a phenol, as is **curcumin**, a yellow pigment isolated from turmeric.
- Many **phenols are antioxidants**, compounds that prevent unwanted oxidation reactions from occurring. Two examples are naturally occurring **vitamin E** and **synthetic BHT**. The OH group on the benzene ring is the key functional group that prevents oxidation reactions from taking place.



vitamin E



BHT
(butylated hydroxy toluene)

End